

CAUSAL REASONING WITH CYCLIC GRAPHS

SIMON FERREIRA

simon.ferreira@sorbonne-universite.fr

L'INSTITUT PIERRE LOUIS D'ÉPIDÉMIOLOGIE ET DE SANTÉ
PUBLIQUE, INSERM, SORBONNE UNIVERSITÉ

JUNE, 2025




- 1 The SCM framework: reminder of Monday
- 2 Cluster DMGs over ADMGs
 - Definitions
 - An example: Summary causal graphs
 - Causal tools in C-DMGs over ADMGs
- 3 The Input/Output SCM framework
 - Definitions
 - Causal tools in DMGs
- 4 Cluster DMGs over DMGs
 - Definitions
 - Causal tools in C-DMGs over DMGs
- 5 Conclusion


1

**THE SCM FRAMEWORK: REMINDER OF
MONDAY**


CAUSAL INFERENCE AND IDENTIFIABILITY: REMINDER OF MONDAY



CAUSAL INFERENCE AND IDENTIFIABILITY: REMINDER OF MONDAY



CAUSAL INFERENCE AND IDENTIFIABILITY: REMINDER OF MONDAY



The causal effect is said to be **identifiable** if it is uniquely computable from $P(\mathbb{V})$.

TOTAL EFFECT A.K.A AVERAGE TREATMENT EFFECT (ATE): REMINDER OF MONDAY

Total effect

$$\begin{aligned} &= \mathbb{E}(Y \mid \text{do}(X = x)) - \mathbb{E}(Y \mid \text{do}(X = x')) \\ &\sim P(y \mid \text{do}(x)) \end{aligned}$$

Where $\text{do}(\cdot)$ is an operator representing an intervention.

Pearl, “Causal diagrams for empirical research”. Biometrika, 1995


TOTAL EFFECT A.K.A AVERAGE TREATMENT EFFECT (ATE): REMINDER OF MONDAY

Total effect

$$= \mathbb{E}(Y \mid \text{do}(X = x)) - \mathbb{E}(Y \mid \text{do}(X = x'))$$

$$\sim P(y \mid \text{do}(x))$$

Where $\text{do}(\cdot)$ is an operator representing an intervention.



$P(y \mid \text{do}(x))$ is **identifiable** if it is uniquely computable from a **positive observational** distribution $P(\mathbb{V})$.

Pearl, "Causal diagrams for empirical research". Biometrika, 1995


TOTAL EFFECT A.K.A AVERAGE TREATMENT EFFECT (ATE): REMINDER OF MONDAY

Total effect

$$= \mathbb{E}(Y \mid \text{do}(X = x)) - \mathbb{E}(Y \mid \text{do}(X = x'))$$

$$\sim P(y \mid \text{do}(x))$$

Where $\text{do}(\cdot)$ is an operator representing an intervention.



$P(y \mid \text{do}(x))$ is **identifiable** if it is uniquely computable from a **positive observational** distribution $P(\mathbb{V})$.

The **do-calculus** is a **sound and complete** for identification.

Pearl, “Causal diagrams for empirical research”. Biometrika, 1995

Structural Causal Model

$$\forall x, \xi_x \leftarrow \mathcal{D}_x$$

$$A := f_A(\xi_a, \xi_{ab})$$

$$G := f_G(\xi_g)$$

$$H := f_H(G, \xi_h)$$

$$I := f_I(G, \xi_i)$$

$$B := f_B(A, H, \xi_b, \xi_{ab})$$

$$C := f_C(A, B, I, \xi_c)$$

$$F := f_F(C, G, \xi_f)$$

$$D := f_D(C, F, \xi_d)$$

$$E := f_E(B, D, G, \xi_e)$$


Pearl, Causality: Models, Reasoning, and Inference. Cambridge University Press, 2009

CAUSAL GRAPHS: REMINDER OF MONDAY

Structural Causal Model

$$\begin{aligned} \forall x, \xi_x &\leftarrow \mathcal{D}_x \\ A &:= f_A(\xi_a, \xi_{ab}) \\ G &:= f_G(\xi_g) \\ H &:= f_H(G, \xi_h) \\ I &:= f_I(G, \xi_i) \\ B &:= f_B(A, H, \xi_b, \xi_{ab}) \\ C &:= f_C(A, B, I, \xi_c) \\ F &:= f_F(C, G, \xi_f) \\ D &:= f_D(C, F, \xi_d) \\ E &:= f_E(B, D, G, \xi_e) \end{aligned}$$

Directed Acyclic Graph (DAG)




Pearl, Causality: Models, Reasoning, and Inference. Cambridge University Press, 2009

CAUSAL GRAPHS: REMINDER OF MONDAY


Structural Causal Model

$$\begin{aligned} \forall x, \xi_x &\leftarrow \mathcal{D}_x \\ A &:= f_A(\xi_a, \xi_{ab}) \\ G &:= f_G(\xi_g) \\ H &:= f_H(G, \xi_h) \\ I &:= f_I(G, \xi_i) \\ B &:= f_B(A, H, \xi_b, \xi_{ab}) \\ C &:= f_C(A, B, I, \xi_c) \\ F &:= f_F(C, G, \xi_f) \\ D &:= f_D(C, F, \xi_d) \\ E &:= f_E(B, D, G, \xi_e) \end{aligned}$$

Directed Acyclic Graph (DAG)



Acyclic Directed Mixed Graph (ADMG)




Pearl, Causality: Models, Reasoning, and Inference. Cambridge University Press, 2009

CAUSAL GRAPHS: REMINDER OF MONDAY


Structural Causal Model

$\forall X, \xi_X \leftarrow \mathcal{D}_X$
 $A := f_A(\xi_a, \xi_{ab})$
 $G := f_G(\xi_g)$
 $H := f_H(G, \xi_h)$
 $I := f_I(G, \xi_i)$
 $B := b$
 $C := f_C(A, B, I, \xi_c)$
 $F := f_F(C, G, \xi_f)$
 $D := f_D(C, F, \xi_d)$
 $E := f_E(B, D, G, \xi_e)$

Directed Acyclic Graph (DAG)



Acyclic Directed Mixed Graph (ADMG)



Pearl, Causality: Models, Reasoning, and Inference. Cambridge University Press, 2009

A path is said to be **blocked** by a set of vertices $\mathbb{Z} \subset \mathbb{V}$ if:

- it contains a chain $\langle A \ast \rightarrow B \rightarrow C \rangle$ or $\langle A \leftarrow B \leftarrow \ast C \rangle$ or a fork $\langle A \leftarrow B \rightarrow C \rangle$ and $B \in \mathbb{Z}$; or
- it contains a collider $\langle A \ast \rightarrow B \leftarrow \ast C \rangle$ such that no descendant of B is in \mathbb{Z} .

Pearl, Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference. Morgan Kaufmann Publishers Inc. 1988

A path is said to be **blocked** by a set of vertices $\mathbb{Z} \subset \mathbb{V}$ if:

- it contains a chain $\langle A \ast \rightarrow B \rightarrow C \rangle$ or $\langle A \leftarrow B \leftarrow \ast C \rangle$ or a fork $\langle A \leftarrow B \rightarrow C \rangle$ and $B \in \mathbb{Z}$; or
- it contains a collider $\langle A \ast \rightarrow B \leftarrow \ast C \rangle$ such that no descendant of B is in \mathbb{Z} .

\mathbb{X} and \mathbb{Y} are **d-separated** by \mathbb{Z} if every path between \mathbb{X} and \mathbb{Y} is blocked by \mathbb{Z} and we write $(\mathbb{X} \perp\!\!\!\perp_d \mathbb{Y} \mid \mathbb{Z})_{\mathcal{G}}$.

Pearl, Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference. Morgan Kaufmann Publishers Inc. 1988

A path is said to be **blocked** by a set of vertices $\mathbb{Z} \subset \mathbb{V}$ if:

- it contains a chain $\langle A \ast \rightarrow B \rightarrow C \rangle$ or $\langle A \leftarrow B \leftarrow \ast C \rangle$ or a fork $\langle A \leftarrow B \rightarrow C \rangle$ and $B \in \mathbb{Z}$; or
- it contains a collider $\langle A \ast \rightarrow B \leftarrow \ast C \rangle$ such that no descendant of B is in \mathbb{Z} .

\mathbb{X} and \mathbb{Y} are **d-separated** by \mathbb{Z} if every path between \mathbb{X} and \mathbb{Y} is blocked by \mathbb{Z} and we write $(\mathbb{X} \perp\!\!\!\perp_d \mathbb{Y} \mid \mathbb{Z})_{\mathcal{G}}$.

Theorem

$$(\mathbb{X} \perp\!\!\!\perp_d \mathbb{Y} \mid \mathbb{Z})_{\mathcal{G}} \Rightarrow \mathbb{X} \perp\!\!\!\perp_{\text{Pr}} \mathbb{Y} \mid \mathbb{Z}$$

Pearl, Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference. Morgan Kaufmann Publishers Inc. 1988

The **do-calculus** consists of three rules:

Rule 1 $P(y|do(x), z, w) = P(y|do(x), w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{X}}}$

Rule 2 $P(y|do(x, z), w) = P(y|do(x), z, w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{XZ}}}$

Rule 3 $P(y|do(x, z), w) = P(y|do(x), w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{XZ(W)}}}$

Pearl, "Causal diagrams for empirical research". Biometrika, 1995

The **do-calculus** consists of three rules:

Rule 1 $P(y|\text{do}(x), z, w) = P(y|\text{do}(x), w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{X}}}$

Rule 2 $P(y|\text{do}(x, z), w) = P(y|\text{do}(x), z, w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{XZ}}}$

Rule 3 $P(y|\text{do}(x, z), w) = P(y|\text{do}(x), w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{XZ(W)}}}$

Theorem

$P(y \mid \text{do}(x))$ is identifiable if and only if there exists a finite sequence of transformations, each conforming to either one of the Rules 1-3 or some standard probability manipulations, that reduces $P(y \mid \text{do}(x))$ into a do-free formula.

Pearl, "Causal diagrams for empirical research". Biometrika, 1995

The **do-calculus** consists of three rules:

Rule 1 $P(y|\text{do}(x), z, w) = P(y|\text{do}(x), w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{X}}}$

Rule 2 $P(y|\text{do}(x, z), w) = P(y|\text{do}(x), z, w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{XZ}}}$

Rule 3 $P(y|\text{do}(x, z), w) = P(y|\text{do}(x), w)$ if $(Y \perp\!\!\!\perp_d Z \mid X, W)_{\mathcal{G}_{\overline{XZ(W)}}}$

Theorem

$P(y \mid \text{do}(x))$ is identifiable if and only if there exists a finite sequence of transformations, each conforming to either one of the Rules 1-3 or some standard probability manipulations, that reduces $P(y \mid \text{do}(x))$ into a do-free formula.

The do-calculus is sound and complete!

Pearl, "Causal diagrams for empirical research". Biometrika, 1995

Definition

A set of variables \mathbb{Z} satisfies the back-door criterion relative to (X, Y) if:

- $\mathbb{Z} \cap \mathbf{De}(X, \mathcal{G}) = \emptyset$, and
- \mathbb{Z} blocks every back-door path (i.e., $\langle X \leftarrow \cdots Y \rangle$).

Definition

A set of variables \mathbb{Z} satisfies the back-door criterion relative to (X, Y) if:

- $\mathbb{Z} \cap \mathbf{De}(X, \mathcal{G}) = \emptyset$, and
- \mathbb{Z} blocks every back-door path (i.e., $\langle X \leftarrow \cdots Y \rangle$).

Theorem


If \mathbb{Z} satisfies the back-door criterion relative to (X, Y) then:

$$\Pr(y \mid do(x)) = \sum_{\mathbb{Z}} \Pr(y \mid x, \mathbb{Z}) \Pr(\mathbb{Z})$$

Definition


A set of variables \mathbb{Z} satisfies the back-door criterion relative to (X, Y) if:

- $\mathbb{Z} \cap \mathbf{De}(X, \mathcal{G}) = \emptyset$, and
- \mathbb{Z} blocks every back-door path (i.e., $\langle X \leftarrow \dots Y \rangle$).




$$\begin{aligned}
 &P(y \mid do(x)) \\
 &= \sum_z P(y \mid do(x), z) P(z \mid do(x)) \\
 &= \sum_z P(y \mid x, z) P(z \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_z P(y \mid x, z) P(z) \quad (\text{Rule 3})
 \end{aligned}$$

THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY




THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY

$$P(y \mid do(x))$$




THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY

$$P(y \mid do(x)) = \sum_w P(y \mid do(x), w) P(w \mid do(x))$$




THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY

$$\begin{aligned}
 P(y \mid do(x)) &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid do(x)) \quad (\text{Rule 2})
 \end{aligned}$$




THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY




$$\begin{aligned}
 P(y \mid do(x)) &= \sum_{w} P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_{w} P(y \mid do(x), do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_{w} P(y \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 3})
 \end{aligned}$$

THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY




$$\begin{aligned}
 P(y \mid do(x)) &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_w P(y \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 3}) \\
 &= \sum_{w, x'} P(y \mid do(w), x') P(x' \mid do(w)) P(w \mid do(x))
 \end{aligned}$$

THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY




$$\begin{aligned}
 P(y \mid do(x)) &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_w P(y \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 3}) \\
 &= \sum_{w, x'} P(y \mid do(w), x') P(x' \mid do(w)) P(w \mid do(x)) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x' \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 2})
 \end{aligned}$$

THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY




$$\begin{aligned}
 P(y \mid do(x)) &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_w P(y \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 3}) \\
 &= \sum_{w, x'} P(y \mid do(w), x') P(x' \mid do(w)) P(w \mid do(x)) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x' \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x') P(w \mid do(x)) \quad (\text{Rule 3})
 \end{aligned}$$

THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY



$$\begin{aligned}
 P(y \mid do(x)) &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_w P(y \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 3}) \\
 &= \sum_{w, x'} P(y \mid do(w), x') P(x' \mid do(w)) P(w \mid do(x)) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x' \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x') P(w \mid do(x)) \quad (\text{Rule 3}) \\
 &= \dots
 \end{aligned}$$

THE DO-CALCULUS CAN TIME BE CONSUMING: REMINDER OF MONDAY



$$\begin{aligned}
 P(y \mid do(x)) &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_w P(y \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 3}) \\
 &= \sum_{w, x'} P(y \mid do(w), x') P(x' \mid do(w)) P(w \mid do(x)) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x' \mid do(w)) P(w \mid do(x)) \quad (\text{Rule 2}) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x') P(w \mid do(x)) \quad (\text{Rule 3}) \\
 &= \dots
 \end{aligned}$$

In this specific case, the total effect is not identifiable
 \implies We can never find a do-free formula!

2

CLUSTER DMGS OVER ADMGS

DEFINITIONS

AN EXAMPLE: SUMMARY CAUSAL GRAPHS

CAUSAL TOOLS IN C-DMGS OVER ADMGS

2

CLUSTER DMGS OVER ADMGS


DEFINITIONS

AN EXAMPLE: SUMMARY CAUSAL GRAPHS

CAUSAL TOOLS IN C-DMGS OVER ADMGS

ADMG


C-DMG over ADMGs



Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs.

ADMG


C-DMG over ADMGs



Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs.


ADMG

C-DMG over ADMGs




Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs.

ADMG




C-DMG over ADMGs




Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs.

ADMG




C-DMG over ADMGs




CYCLES !!!!

Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs.

ADMG



C-DMG over ADMGs



Assumption: For every cycle (e.g., $(B, C) \rightarrow (D, E, F, G) \rightarrow (H) \rightarrow (B, C)$) no 2 adjacent clusters are of size 1.

Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs.


2

CLUSTER DMGS OVER ADMGS


DEFINITIONS

AN EXAMPLE: SUMMARY CAUSAL GRAPHS


CAUSAL TOOLS IN C-DMGS OVER ADMGS



Assaad, Devijver, and Gaussier, “Survey and Evaluation of Causal Discovery Methods for Time Series”. JAIR, 2022




Summary causal graph (SCG)




Assaad, Devijver, and Gaussier, “Survey and Evaluation of Causal Discovery Methods for Time Series”. JAIR, 2022

ADMGs WITH TIME AND SUMMARY CAUSAL GRAPHS




Summary causal graph (SCG)




■ SCGs can contain cycles

Assaad, Devijver, and Gaussier, “Survey and Evaluation of Causal Discovery Methods for Time Series”. JAIR, 2022




Summary causal graph (SCG)




- SCGs can contain cycles
- Each vertex does not map to one single random variable

Assaad, Devijver, and Gaussier, “Survey and Evaluation of Causal Discovery Methods for Time Series”. JAIR, 2022



Summary causal graph (SCG)



- SCGs can contain cycles
- Each vertex does not map to one single random variable
- There might exist many ADMGs compatible with one SCG

Assaad, Devijver, and Gaussier, “Survey and Evaluation of Causal Discovery Methods for Time Series”. JAIR, 2022

TWO TYPES OF TOTAL EFFECTS IN C-DMGS


A micro total effect is a total effect from a single variable in a cluster (e.g., $X_{t-\gamma}$) to another single variable (e.g., Y_t). For example: $\Pr(y_t \mid \text{do}(x_{t-\gamma}))$.

Ferreira and Assaad, “Identifying macro conditional independencies and macro total effects in summary causal graphs with latent confounding”

An example: Summary causal graphs

TWO TYPES OF TOTAL EFFECTS IN C-DMGS

A micro total effect is a total effect from a single variable in a cluster (e.g., $X_{t-\gamma}$) to another single variable (e.g., Y_t). For example: $\Pr(y_t \mid \text{do}(x_{t-\gamma}))$.




Ferreira and Assaad, “Identifying macro conditional independencies and macro total effects in summary causal graphs with latent confounding”

An example: Summary causal graphs

TWO TYPES OF TOTAL EFFECTS IN C-DMGS

A micro total effect is a total effect from a single variable in a cluster (e.g., $X_{t-\gamma}$) to another single variable (e.g., Y_t). For example: $\Pr(y_t \mid \text{do}(x_{t-\gamma}))$.




A macro total effect is a total effect from a whole cluster (e.g., $\{X_{t_0}, X_{t_0+1}, \dots, X_{t-1}, X_t\}$) to a whole other cluster (e.g., $\{Y_{t_0}, Y_{t_0+1}, \dots, Y_{t-1}, Y_t\}$). For example: $\Pr(\{y_{t_0}, y_{t_0+1}, \dots, y_{t-1}, y_t\} \mid \text{do}(\{x_{t_0}, x_{t_0+1}, \dots, x_{t-1}, x_t\}))$.

Ferreira and Assaad, “Identifying macro conditional independencies and macro total effects in summary causal graphs with latent confounding”


An example: Summary causal graphs

TWO TYPES OF TOTAL EFFECTS IN C-DMGS

A micro total effect is a total effect from a single variable in a cluster (e.g., $X_{t-\gamma}$) to another single variable (e.g., Y_t). For example: $\Pr(y_t \mid \text{do}(x_{t-\gamma}))$.



A macro total effect is a total effect from a whole cluster (e.g., $\{X_{t_0}, X_{t_0+1}, \dots, X_{t-1}, X_t\}$) to a whole other cluster (e.g., $\{Y_{t_0}, Y_{t_0+1}, \dots, Y_{t-1}, Y_t\}$). For example: $\Pr(\{y_{t_0}, y_{t_0+1}, \dots, y_{t-1}, y_t\} \mid \text{do}(\{x_{t_0}, x_{t_0+1}, \dots, x_{t-1}, x_t\}))$.




Ferreira and Assaad, "Identifying macro conditional independencies and macro total effects in summary causal graphs with latent confounding".


An example: Summary causal graphs

TWO TYPES OF TOTAL EFFECTS IN C-DMGS

A micro total effect is a total effect from a single variable in a cluster (e.g., $X_{t-\gamma}$) to another single variable (e.g., Y_t). For example: $\Pr(Y_t \mid \text{do}(X_{t-\gamma}))$.




A macro total effect is a total effect from a whole cluster (e.g., $\{X_{t_0}, X_{t_0+1}, \dots, X_{t-1}, X_t\}$) to a whole other cluster (e.g., $\{Y_{t_0}, Y_{t_0+1}, \dots, Y_{t-1}, Y_t\}$). For example: $\Pr(\{Y_{t_0}, Y_{t_0+1}, \dots, Y_{t-1}, Y_t\} \mid \text{do}(\{X_{t_0}, X_{t_0+1}, \dots, X_{t-1}, X_t\}))$.



Ferreira and Assaad, "Identifying macro conditional independencies and macro total effects in summary causal graphs with latent confounding".

An example: Summary causal graphs



2

CLUSTER DMGS OVER ADMGS

DEFINITIONS

AN EXAMPLE: SUMMARY CAUSAL GRAPHS

CAUSAL TOOLS IN C-DMGS OVER ADMGS

Is d-separation applicable for C-DMGs over ADMGs?

Is d-separation applicable for C-DMGs over ADMGs?

Yes!

Is d-separation applicable for C-DMGs over ADMGs?


Yes!

Theorem


d-separation is valid in C-DMGs over ADMGs.

- *If a d-separation holds in a given C-DMG, then it holds in every compatible ADMG.*
- *If a d-separation does not hold in a given C-DMG, then there exists a compatible ADMG in which it does not hold.*

DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs




DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



$$X \stackrel{?}{\perp\!\!\!\perp}_{\mathcal{G}} Y \mid Z, W$$


DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



Is the path $\langle X \rightarrow Z \rightarrow Y \rangle$
blocked by $\{Z, W\}$?

$$X \stackrel{?}{\perp\!\!\!\perp}_{\mathcal{G}} Y \mid Z, W$$


DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



Is the path $\langle X \leftarrow Z \rightarrow Y \rangle$
blocked by $\{Z, W\}$?

$$X \stackrel{?}{\perp\!\!\!\perp}_{\mathcal{G}} Y \mid Z, W$$


DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



Is the path $\langle X \leftarrow W \rightarrow Y \rangle$
blocked by $\{Z, W\}$?

$$X \stackrel{?}{\perp\!\!\!\perp}_{\mathcal{G}} Y \mid Z, W$$


DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



Is the path $\langle X \leftarrow W \leftarrow Y \rangle$
blocked by $\{Z, W\}$?


$$X \stackrel{?}{\perp\!\!\!\perp}_{\mathcal{G}} Y \mid Z, W$$

DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



$$X \perp\!\!\!\perp_{\mathcal{G}} Y \mid Z, W$$


DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



$$X \perp\!\!\!\perp_{\mathcal{G}} Y \mid Z, W$$

$$W \stackrel{?}{\perp\!\!\!\perp}_{\mathcal{G}} Z \mid \Omega \setminus \{X\}$$

DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs




Is the path $\langle W \rightarrow X \rightarrow Z \rangle$
 blocked by $\Omega \setminus \{X\}$?

$$X \perp\!\!\!\perp_{\mathcal{G}} Y \mid Z, W$$

$$W \stackrel{?}{\perp\!\!\!\perp}_{\mathcal{G}} Z \mid \Omega \setminus \{X\}$$


DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



$$X \perp\!\!\!\perp_Y Y \mid Z, W$$

$$W \not\perp\!\!\!\perp_Y Z \mid \Omega \setminus \{X\}$$

DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs




$$X \perp\!\!\!\perp_{\mathcal{G}} Y \mid Z, W$$

$$W \not\perp\!\!\!\perp_{\mathcal{G}} Z \mid \Omega \setminus \{X\}$$

$$W \stackrel{?}{\perp\!\!\!\perp}_{\mathcal{G}} Z \mid \Omega \cup \{X\}$$

DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs




Is the path $\langle W \rightarrow X \leftarrow Z \rangle$
 blocked by $\Omega \cup \{X\}$?

$$X \perp\!\!\!\perp_Y Y \mid Z, W$$

$$W \not\perp\!\!\!\perp_Z Z \mid \Omega \setminus \{X\}$$

$$W \stackrel{?}{\perp\!\!\!\perp}_Z Z \mid \Omega \cup \{X\}$$

DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs




$$X \perp\!\!\!\perp_Y Y \mid Z, W$$

$$W \not\perp\!\!\!\perp_Y Z \mid \Omega \setminus \{X\}$$

$$W \not\perp\!\!\!\perp_Y Z \mid \Omega \cup \{X\}$$

DEMO: D-SEPARATION FOR C-DMGs OVER ADMGs



$$X \perp\!\!\!\perp_{\mathcal{G}} Y \mid Z, W$$

$$W \not\perp\!\!\!\perp_{\mathcal{G}} Z \mid \Omega \setminus \{X\}$$

$$W \not\perp\!\!\!\perp_{\mathcal{G}} Z \mid \Omega \cup \{X\}$$

$$X \perp\!\!\!\perp_{\mathcal{G}} Y \mid Z, W$$

Is do-calculus applicable for C-DMGs over ADMGs?

Is do-calculus applicable for C-DMGs over ADMGs?

Yes!

Is do-calculus applicable for C-DMGs over ADMGs?


Yes!

Theorem


The Rules 1-3 of the do-calculus are valid in C-DMGs over DMGs.

- *If a sequence of rules apply in a given C-DMG, then it applies in every compatible ADMG.*
- *If a sequence of rules of the do-calculus does not apply in a given C-DMG, then there exists a compatible ADMG in which it does not apply.*

DEMO: DO-CALCULUS FOR C-DMGs OVER ADMGs





DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGs





$$\begin{aligned}
 P(y \mid do(x)) \\
 &= P(y \mid \cancel{x}) \quad (\text{Rule 2})
 \end{aligned}$$

DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGs



$$\begin{aligned}
 &P(y \mid do(x)) \\
 &= P(y \mid \textcolor{red}{x}) \quad (\text{Rule 2})
 \end{aligned}$$

DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGS





$$P(y \mid do(x))$$

$$P(y \mid do(x))$$

$$= P(y \mid \cancel{x}) \quad (\text{Rule 2})$$



DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGs



$$\begin{aligned}
 &P(y \mid do(x)) \\
 &= \sum_{w} P(y \mid do(x), w) P(w \mid do(x))
 \end{aligned}$$

$$\begin{aligned}
 &P(y \mid do(x)) \\
 &= P(y \mid x) \quad (\text{Rule 2})
 \end{aligned}$$



DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGs



$$\begin{aligned}
 &P(y \mid do(x)) \\
 &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid x) \quad (\text{Rule 2}) \times 2
 \end{aligned}$$

$$\begin{aligned}
 &P(y \mid do(x)) \\
 &= P(y \mid x) \quad (\text{Rule 2})
 \end{aligned}$$



DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGS



$$\begin{aligned}
 P(y \mid do(x)) \\
 &= P(y \mid x) \quad (\text{Rule 2})
 \end{aligned}$$

$$\begin{aligned}
 P(y \mid do(x)) \\
 &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid x) \quad (\text{Rule 2}) \times 2 \\
 &= \sum_w P(y \mid do(w)) P(w \mid x) \quad (\text{Rule 3})
 \end{aligned}$$



DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGS



$$\begin{aligned}
 P(y \mid do(x)) \\
 &= P(y \mid x) \quad (\text{Rule 2})
 \end{aligned}$$

$$\begin{aligned}
 &P(y \mid do(x)) \\
 &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid x) \quad (\text{Rule 2}) \times 2 \\
 &= \sum_w P(y \mid do(w)) P(w \mid x) \quad (\text{Rule 3}) \\
 &= \sum_{w, x'} P(y \mid do(w), x') P(x' \mid do(w)) P(w \mid x)
 \end{aligned}$$



DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGS



$$\begin{aligned}
 P(y \mid do(x)) \\
 &= P(y \mid x) \quad (\text{Rule 2})
 \end{aligned}$$

$$\begin{aligned}
 &P(y \mid do(x)) \\
 &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid x) \quad (\text{Rule 2}) \times 2 \\
 &= \sum_w P(y \mid do(w)) P(w \mid x) \quad (\text{Rule 3}) \\
 &= \sum_{w, x'} P(y \mid do(w), x') P(x' \mid do(w)) P(w \mid x) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x' \mid do(w)) P(w \mid x) \quad (\text{Rule 2})
 \end{aligned}$$

DEMO: DO-CALCULUS FOR C-DMGS OVER ADMGS



$$\begin{aligned}
 P(y \mid do(x)) \\
 &= P(y \mid x) \quad (\text{Rule 2})
 \end{aligned}$$

$$\begin{aligned}
 P(y \mid do(x)) \\
 &= \sum_w P(y \mid do(x), w) P(w \mid do(x)) \\
 &= \sum_w P(y \mid do(x), do(w)) P(w \mid x) \quad (\text{Rule 2}) \times 2 \\
 &= \sum_w P(y \mid do(w)) P(w \mid x) \quad (\text{Rule 3}) \\
 &= \sum_{w, x'} P(y \mid do(w), x') P(x' \mid do(w)) P(w \mid x) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x' \mid do(w)) P(w \mid x) \quad (\text{Rule 2}) \\
 &= \sum_{w, x'} P(y \mid w, x') P(x') P(w \mid x) \quad (\text{Rule 3})
 \end{aligned}$$

Reminder:

- A macro causal effect is a causal effect from a whole cluster \mathbb{X} to a another whole cluster \mathbb{Y} , e.g.,
 $\Pr(Y_{t_0}, Y_{t_0+1}, \dots, Y_{t-1}, Y_t \mid \text{do}(X_{t_0}, X_{t_0+1}, \dots, X_{t-1}, X_t)).$
- A micro causal effect is a total effect from a single variable X to another single variable Y , e.g., $\Pr(y_t \mid \text{do}(x_{t-\gamma}))$.

Reminder:

- A macro causal effect is a causal effect from a whole cluster \mathbb{X} to another whole cluster \mathbb{Y} , e.g.,
 $\Pr(Y_{t_0}, Y_{t_0+1}, \dots, Y_{t-1}, Y_t \mid \text{do}(X_{t_0}, X_{t_0+1}, \dots, X_{t-1}, X_t)).$
- A micro causal effect is a total effect from a single variable X to another single variable Y , e.g., $\Pr(y_t \mid \text{do}(x_{t-\gamma}))$.

Can we identify micro causal effects in C-DMGs over ADMGs?

Reminder:

- A macro causal effect is a causal effect from a whole cluster \mathbb{X} to another whole cluster \mathbb{Y} , e.g.,
 $\Pr(Y_{t_0}, Y_{t_0+1}, \dots, Y_{t-1}, Y_t \mid \text{do}(X_{t_0}, X_{t_0+1}, \dots, X_{t-1}, X_t)).$
- A micro causal effect is a total effect from a single variable X to another single variable Y , e.g., $\Pr(y_t \mid \text{do}(x_{t-\gamma}))$.

Can we identify micro causal effects in C-DMGs over ADMGs?

In specific cases and/or with prior knowledge, such as time,
then **yes!**

Can we identify micro causal effects in C-DMGs over ADMGs?

In specific cases and/or with prior knowledge, such as time,
then **yes!**

- Assaad et al., “Identifiability of total effects from abstractions of time series causal graphs”.
- Assaad, “Towards identifiability of micro total effects in summary causal graphs with latent confounding: extension of the front-door criterion”.
- Ferreira and Assaad, “Identifiability of Direct Effects from Summary Causal Graphs”.
- Ferreira and Assaad, “Average Controlled and Average Natural Micro Direct Effects in Summary Causal Graphs”.

3

THE INPUT/OUTPUT SCM FRAME- WORK

DEFINITIONS

CAUSAL TOOLS IN DMGs

3

THE INPUT/OUTPUT SCM FRAME- WORK

DEFINITIONS

CAUSAL TOOLS IN DMGS

i/o SCM

$$\forall x, \xi_x \leftarrow \mathcal{D}_x$$

$$A := f_A(\xi_a, \xi_{ab})$$

$$B := f_B(A, D, E, \xi_b, \xi_{ab})$$

$$C := f_C(B, \xi_c)$$

$$D := f_D(C, \xi_d)$$

$$E := f_E(C, \xi_e)$$

$$(B, C, D) := f_{(B,C,D)}(A, E, \xi_b, \xi_{ab}, \xi_c, \xi_d)$$

$$(B, C, E) := f_{(B,C,E)}(A, D, \xi_b, \xi_{ab}, \xi_c, \xi_e)$$

$$(B, C, D, E) := f_{(B,C,D)}(A, \xi_b, \xi_{ab}, \xi_c, \xi_d, \xi_e)$$

Assumption: Compatibility of the generating processes in cycles.

$$\text{e.g., } (b, c, d) = f_{(B,C,D)}(a, e, \xi_b, \xi_{ab}, \xi_c, \xi_d) \implies b = f_B(a, d, e, \xi_b, \xi_{ab})$$

i/o SCM

$$\forall X, \xi_X \leftarrow \mathcal{D}_X$$

$$A := f_A(\xi_a, \xi_{ab})$$

$$B := f_B(A, D, E, \xi_b, \xi_{ab})$$

$$C := f_C(B, \xi_c)$$

$$D := f_D(C, \xi_d)$$


$$E := f_E(C, \xi_e)$$

$$(B, C, D) := f_{(B,C,D)}(A, E, \xi_b, \xi_{ab}, \xi_c, \xi_d)$$

$$(B, C, E) := f_{(B,C,E)}(A, D, \xi_b, \xi_{ab}, \xi_c, \xi_e)$$

$$(B, C, D, E) := f_{(B,C,D)}(A, \xi_b, \xi_{ab}, \xi_c, \xi_d, \xi_e)$$

Acyclic Directed Mixed Graph (DMG)



Forré and Mooij, “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”

i/o SCM

$$\forall \mathbf{x}, \xi_{\mathbf{x}} \leftarrow \mathcal{D}_{\mathbf{x}}$$

$$A := f_A(\xi_a, \xi_{ab})$$

$$B := f_B(A, D, E, \xi_b, \xi_{ab})$$

$$C := f_C(B, \xi_c)$$

$$D := d$$


$$E := f_E(C, \xi_e)$$

~~$$(B, C, D) := f_{(B,C,D)}(A, E, \xi_b, \xi_{ab}, \xi_c, \xi_d)$$~~

$$(B, C, E) := f_{(B,C,E)}(A, D, \xi_b, \xi_{ab}, \xi_c, \xi_e)$$

~~$$(B, C, D, E) := f_{(B,C,D)}(A, \xi_b, \xi_{ab}, \xi_c, \xi_d, \xi_e)$$~~

Directed Mixed Graph (DMG)



Forré and Mooij, “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”

3

THE INPUT/OUTPUT SCM FRAME- WORK

DEFINITIONS

CAUSAL TOOLS IN DMGs

Strongly connected component: $Scc(V, \mathcal{G}) := An(V, \mathcal{G}) \cap De(V, \mathcal{G})$

A walk $\tilde{\pi} = \langle V_1, \dots, V_n \rangle$ is said to be σ -blocked by a set of vertices $\mathbb{Z} \subseteq \mathbb{V}$ if:

1. $\exists 1 < i < n$ such that $\langle V_{i-1} \xrightarrow{*} V_i \xleftarrow{*} V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \notin \mathbb{Z}$, or
2. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftarrow V_i \xleftarrow{*} V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus Scc(V_{i-1}, \mathcal{G})$, or
3. $\exists 1 < i < n$ such that $\langle V_{i-1} \xrightarrow{*} V_i \rightarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus Scc(V_{i+1}, \mathcal{G})$, or
4. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftarrow V_i \rightarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus (Scc(V_{i-1}, \mathcal{G}) \cap Scc(V_{i+1}, \mathcal{G}))$.

Forré and Mooij, “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”. PMLR, 2020.

Strongly connected component: $Scc(V, \mathcal{G}) := An(V, \mathcal{G}) \cap De(V, \mathcal{G})$

A walk $\tilde{\pi} = \langle V_1, \dots, V_n \rangle$ is said to be **σ -blocked** by a set of vertices $\mathbb{Z} \subseteq \mathbb{V}$ if:

1. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftrightarrow V_i \leftrightarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \notin \mathbb{Z}$, or
2. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftarrow V_i \leftarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus Scc(V_{i-1}, \mathcal{G})$, or
3. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftrightarrow V_i \rightarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus Scc(V_{i+1}, \mathcal{G})$, or
4. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftarrow V_i \rightarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus (Scc(V_{i-1}, \mathcal{G}) \cap Scc(V_{i+1}, \mathcal{G}))$.

\mathbb{X} and \mathbb{Y} are **σ -separated** by \mathbb{Z} if every **walk** between \mathbb{X} and \mathbb{Y} is blocked by \mathbb{Z} and we write $(\mathbb{X} \perp\!\!\!\perp_{\sigma} \mathbb{Y} \mid \mathbb{Z})_{\mathcal{G}}$.

Forré and Mooij, “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”. PMLR, 2020.

Strongly connected component: $Scc(V, \mathcal{G}) := An(V, \mathcal{G}) \cap De(V, \mathcal{G})$

A walk $\tilde{\pi} = \langle V_1, \dots, V_n \rangle$ is said to be σ -blocked by a set of vertices $\mathbb{Z} \subseteq \mathbb{V}$ if:

1. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftrightarrow V_i \leftrightarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \notin \mathbb{Z}$, or
2. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftarrow V_i \leftarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus Scc(V_{i-1}, \mathcal{G})$, or
3. $\exists 1 < i < n$ such that $\langle V_{i-1} \rightarrow V_i \rightarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus Scc(V_{i+1}, \mathcal{G})$, or
4. $\exists 1 < i < n$ such that $\langle V_{i-1} \leftarrow V_i \rightarrow V_{i+1} \rangle \subseteq \tilde{\pi}$ and $V_i \in \mathbb{Z} \setminus (Scc(V_{i-1}, \mathcal{G}) \cap Scc(V_{i+1}, \mathcal{G}))$.

\mathbb{X} and \mathbb{Y} are σ -separated by \mathbb{Z} if every walk between \mathbb{X} and \mathbb{Y} is blocked by \mathbb{Z} and we write $(\mathbb{X} \perp\!\!\!\perp_{\sigma} \mathbb{Y} \mid \mathbb{Z})_{\mathcal{G}}$.

Theorem

$$(\mathbb{X} \perp\!\!\!\perp_{\sigma} \mathbb{Y} \mid \mathbb{Z})_{\mathcal{G}} \Rightarrow \mathbb{X} \perp\!\!\!\perp_{Pr} \mathbb{Y} \mid \mathbb{Z}$$

Forré and Mooij, “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”. PMLR, 2020.

The σ -based **do-calculus** consists of three rules:

Rule 1 $P(y|do(x), z, w) = P(y|do(x), w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{X}}}$

Rule 2 $P(y|do(x, z), w) = P(y|do(x), z, w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{XZ}}}$

Rule 3 $P(y|do(x, z), w) = P(y|do(x), w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{XZ}(w)}}$

Forré and Mooij, “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”. PMLR, 2020.

The σ -based **do-calculus** consists of three rules:

Rule 1 $P(y|\text{do}(x), z, w) = P(y|\text{do}(x), w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{X}}}$

Rule 2 $P(y|\text{do}(x, z), w) = P(y|\text{do}(x), z, w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{XZ}}}$

Rule 3 $P(y|\text{do}(x, z), w) = P(y|\text{do}(x), w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{XZ(W)}}}$

Theorem

$P(y \mid \text{do}(x))$ is identifiable if and only if there exists a finite sequence of transformations, each conforming to either one of the Rules 1-3 or some standard probability manipulations, that reduces $P(y \mid \text{do}(x))$ into a do-free formula.

Forré and Mooij, “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”. PMLR, 2020.

The σ -based **do-calculus** consists of three rules:

Rule 1 $P(y|\text{do}(x), z, w) = P(y|\text{do}(x), w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{X}}}$

Rule 2 $P(y|\text{do}(x, z), w) = P(y|\text{do}(x), z, w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{XZ}}}$

Rule 3 $P(y|\text{do}(x, z), w) = P(y|\text{do}(x), w)$ if $(Y \perp\!\!\!\perp_{\sigma} Z \mid X, W)_{\mathcal{G}_{\overline{XZ(W)}}}$

Theorem

$P(y \mid \text{do}(x))$ is identifiable if and only if there exists a finite sequence of transformations, each conforming to either one of the Rules 1-3 or some standard probability manipulations, that reduces $P(y \mid \text{do}(x))$ into a do-free formula.

The do-calculus is sound!

Forré and Mooij, “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”. PMLR, 2020.

4

CLUSTER DMGS OVER DMGS

DEFINITIONS

CAUSAL TOOLS IN C-DMGS OVER DMGS

4


CLUSTER DMGS OVER DMGS

DEFINITIONS

CAUSAL TOOLS IN C-DMGS OVER DMGS

DMG


C-DMG over DMGs



Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs over DMGs.

DMG


C-DMG over DMGs



Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs over DMGs.


DMG

C-DMG over DMGs




Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs over DMGs.

DMG




C-DMG over DMGs




Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs over DMGs.

DMG



C-DMG over DMGs



When clustering, cycles can:

- "disappear" (e.g., $I \rightarrow J \rightarrow I$) and/or
- "remain" (e.g., $C \rightarrow F \rightarrow D \rightarrow C$) and/or
- "appear" (e.g., $A \rightarrow (B, C) \rightarrow A$).

Ferreira and Assaad, Identifying Macro Causal Effects in C-DMGs over DMGs.

4

CLUSTER DMGS OVER DMGS

DEFINITIONS

CAUSAL TOOLS IN C-DMGS OVER DMGS

Is σ -separation applicable for C-DMGs over DMGs?

Is σ -separation applicable for C-DMGs over DMGs?

Yes!

Is σ -separation applicable for C-DMGs over DMGs?

Yes!

Theorem

σ -separation is valid in C-DMGs over DMGs.

- *If a σ -separation holds in a given C-DMG, then it holds in every compatible DMG.*
- *If a σ -separation does not hold in a given C-DMG, then there exists a compatible DMG in which it does not hold.*

Is σ -based do-calculus applicable for C-DMGs over DMGs?

Is σ -based do-calculus applicable for C-DMGs over DMGs?
Yes!

Is σ -based do-calculus applicable for C-DMGs over DMGs?
Yes!

Theorem

The Rules 1-3 of the do-calculus are valid in C-DMGs over DMGs.

- *If a sequence of rules apply in a given C-DMG, then it applies in every compatible DMG.*
- *If a sequence of rules of the do-calculus does not apply in a given C-DMG, then there exists a compatible DMG in which it does not apply.*

5

CONCLUSION

Use causal graphs !

- Even if the causal graph is not fully specified,
- Even if the causal graph is cyclic.

Now, there exist tools which accomodate these cases.

THANK YOU FOR YOUR ATTENTION!

ANY QUESTIONS?





Assaad, Charles K. “Towards identifiability of micro total effects in summary causal graphs with latent confounding: extension of the front-door criterion”. In: Transactions on Machine Learning Research (2025). ISSN: 2835-8856. URL:

<https://openreview.net/forum?id=5f7YlSKG1l>.



Assaad, Charles K., Emilie Devijver, and Eric Gaussier. “Survey and Evaluation of Causal Discovery Methods for Time Series”. In: 73 (May 2022). ISSN: 1076-9757. DOI: [10.1613/jair.1.13428](https://doi.org/10.1613/jair.1.13428). URL: <https://doi.org/10.1613/jair.1.13428>.



Assaad, Charles K. et al. “Identifiability of total effects from abstractions of time series causal graphs”. In: Proceedings of the Fortieth Conference on Uncertainty in Artificial Intelligence. Ed. by Negar Kiyavash and Joris M. Mooij. Vol. 244. Proceedings of Machine Learning Research. PMLR, July 2024, pp. 173–185.



Ferreira, Simon and Charles K. Assaad. “Average Controlled and Average Natural Micro Direct Effects in Summary Causal Graphs”. In: (2025). Submitted.



— . “Identifiability of Direct Effects from Summary Causal Graphs”. In: Proceedings of the AAAI Conference on Artificial Intelligence 38.18 (Mar. 2024), pp. 20387–20394. DOI: **10.1609/aaai.v38i18.30021**.



Ferreira, Simon and Charles K. Assaad.
Identifying Macro Causal Effects in C-DMGs. 2025. arXiv:
2504.01551 [cs.AI].



— .“Identifying macro conditional independencies and macro total effects in summary causal graphs with latent confounding”. In: (2025). Causal Inference for Time Series Data Workshop at the 40th Conference on Uncertainty in Artificial Intelligence.



Forré, Patrick and Joris M. Mooij. “Causal Calculus in the Presence of Cycles, Latent Confounders and Selection Bias”.

In:

Proceedings of The 35th Uncertainty in Artificial Intelligence Conference

Ed. by Ryan P. Adams and Vibhav Gogate. Vol. 115.

Proceedings of Machine Learning Research. PMLR, 2020, pp. 71–80.



Pearl, Judea. “Causal diagrams for empirical research”. In: Biometrika 82.4 (Dec. 1995), pp. 669–688. ISSN: 0006-3444. DOI: [10.1093/biomet/82.4.669](https://doi.org/10.1093/biomet/82.4.669).



— .Causality: Models, Reasoning, and Inference. Cambridge University Press, 2009.



— .Probabilistic Reasoning in Intelligent Systems: Networks of Pl
San Francisco, CA, USA: Morgan Kaufmann Publishers Inc.,
1988. ISBN: 0-934613-73-7.